

RESEARCH

Open Access



Cryptic intermediate snail host of the liver fluke *Fasciola hepatica* in Africa

Anna Mahulu¹, Catharina Clewing¹, Björn Stelbrink^{1,2}, Fred D. Chibwana^{1,3}, Immaculate Tumwebaze¹, J. Russell Stothard⁴ and Christian Albrecht^{1,5*}

Abstract

Background: Snails such as *Galba truncatula* are hosts for trematode flukes causing fascioliasis, a zoonosis that is a major public health problem. *Galba truncatula* has recently been shown to be a cryptic species complex. African populations of *Galba* spp. are not yet studied using molecular assessments and it is imperative to do so and reconstruct the centre of origin of *Galba* and to understand when and by what means it may have colonized the highlands of Africa and to what extent humans might have been involved in that process.

Methods: Samples from all known sub-ranges throughout Africa and new samples from Europe and Asia were obtained. We used a combination of two mitochondrial (*cox1* and *16S*) and one nuclear (ITS2) markers and phylogenetic, divergence time estimates and phylogeographical methods to determine the identity and biogeographical affinities. We also reconstructed the colonization history including the likely mode of dispersal and tested for the presence of cryptic *Galba* species in Africa.

Results: *Galba truncatula* is restricted to the Palaearctic region of the continent, namely Morocco. All sub-Saharan populations proved to be a distinct species according to the phylogenetic analyses and genetic distance. We propose to use the existing name *Galba mweruensis* (Connolly, 1929) for this species which is morphologically indistinguishable from the other two species hitherto known to occur in northern Africa, i.e. *G. truncatula* and *G. schirazensis*. Sub-tropical Africa has been colonized only once in either the Pliocene and possibly Miocene. Diversification within *G. mweruensis* is dated to the Plio-Pleistocene and thus human-mediated dispersal can be ruled out for the initial colonization of the isolated mountain ranges. There are potentially even more cryptic species in high altitude areas of Africa as outlined by the distinctness of the population found at the top of Mt. Elgon, Uganda.

Conclusions: From a novel genetic inspection of available African material, a hitherto neglected distinct species, *G. mweruensis*, now appears a major host of *F. hepatica* throughout sub-Saharan Africa. A closer examination of trematode parasites hosted by this species is needed in order to understand transmission patterns in highlands throughout eastern and southern Africa. We encourage future studies to inspect other high altitude areas in Africa in light of parasites of either veterinary or medical importance.

Keywords: Fascioliasis, Medical malacology, Cryptic species, *Galba truncatula*, Lymnaeidae, Dispersal, Islands-in-the-sky

Background

Parasitic disease caused by the liver flukes of the genus *Fasciola* affects hundreds of millions of people and livestock worldwide. Collectively, they cause considerable

economic damage. Indeed, fascioliasis, a very debilitating snail-borne disease, is widespread across the globe; however, in the subtropical/cooler regions it is caused by *Fasciola hepatica* [1] whereas in the tropical/warmer regions is caused by *Fasciola gigantica* [2].

To complete the life-cycle, the two species of liver fluke are tied to a variety of intermediate freshwater pulmonate snail hosts of the family Lymnaeidae [3].

*Correspondence: christian.albrecht@allzool.bio.uni-giessen.de

¹ Department of Animal Ecology and Systematics, Justus Liebig University Giessen, Giessen, Germany

Full list of author information is available at the end of the article



Until relatively recently, the taxonomy of snails was consolidated to a single genus *Lymnaea* with remarkable morphological diversity; however, with application of molecular phylogenetics a multi-generic nomenclature has become favoured with *Galba* and *Radix* now used in preference [4]. In Africa, for example, *Galba truncatula* (also known as *Lymnaea truncatula*) is involved in the transmission of *F. hepatica* while *Radix natalensis* is involved in the transmission of *F. gigantica* with any epidemiological cross-over considered to be rare [4]. As an intermediate host of *F. hepatica*, the liver fluke largely responsible for human disease, *G. truncatula* is characterized by its amphibious lifestyle, adaptation to cooler habitats, and its ability to withstand drought events and other harsh environmental conditions in unstable waterbodies [5]. It has been found in high altitudes in South America, where it can reach up to 4100 m [6] and it is thus among the few gastropods reaching extreme habitats on high elevations [7].

The taxonomy of lymnaeid gastropods continues to be debated [4, 8], but recent molecular phylogenetic studies improved the understanding of the evolution of this major freshwater gastropod family [3, 9–11]. The species *G. truncatula* has been treated as belonging to *Lymnaea* and *Fossaria* in North America and is thus a prime example of taxonomic confusion in lymnaeid systematics. *Galba truncatula* as the type-species of the genus is conceived to be mainly a Holarctic species [12], with a wide distribution range throughout North America and Eurasia, where it reaches as far as India [13]. The scattered occurrences in South America have been interpreted as recent introductions [14]. However, the real extent of the distribution of *G. truncatula* on a global scale is potentially masked by the occurrence of cryptic species that are morphologically indistinguishable from *G. truncatula*. Among these species are *Lymnaea cubensis* [15] and *Lymnaea schirazensis*, two species that have been previously confused with *G. truncatula* prior to the introduction of molecular methods of characterisation. Such a confusing situation has important implications to parasite transmission and epidemiology because the cryptic species may differ in their competence for transmission of *F. hepatica*.

Given the importance of these species for veterinary and human parasitology, a number of attempts have been made to identify species based on molecular markers. As a result, a relatively rich record of sequences of several mitochondrial and nuclear molecular markers is available for comparative analyses of material studied recently [3]. On the population level, SNPs [16] and microsatellites have been published [17]. A recent study proposed an easy and inexpensive PCR-based approach to distinguish among three cryptic *Galba* species [15].

Despite the variety of applicable molecular diagnostic markers, there is a significant gap of knowledge about snails referred to as *G. truncatula* on the African continent. Here, the *Galba truncatula*-like snails have a disjunct distribution with four largely isolated sub-ranges: in the mountainous parts of the Maghreb states in northern Africa [18], the highlands of Ethiopia [19], some highland areas in East Africa such as Mt. Elgon [20], Usambara Mt. [21], the Kitulo Plateau [22], the highlands of Lesotho [23], and temperate coastal, i.e. cooler, regions of South Africa [24].

When compared to the other native lymnaeid species in Africa, such as *Radix natalensis* the main host of *Fasciola gigantica*, the distribution pattern of what is considered *G. truncatula* is particularly striking (Fig. 1) being confined to allopatry in higher altitudes [20]. The discontinuous range of *G. truncatula* has been hypothesized to be the result of passive dispersal by migratory birds, being more likely perhaps than an alternative of much longer historical associations with geological vicariance of uplifted African highlands [25]. Given scattered subfossil records in the Sahara, the Near East and Namibia [21], this could represent a range of ancestral or relic habitats isolated for eons. Another possibility would be a human- or livestock-mediated introduction, given the well-recognized anthropophily of the species [26]. In fact, historical records in the eastern part of the DR Congo have been attributed to human introductions [13]. Records of the Nile Delta in Egypt recently turned out to represent populations of *Lymnaea schirazensis* [27] and thus raise questions as to a potential camouflaged invasion in other parts of the continent. The only populations of *Galba* spp. that were identified by molecular DNA to be *G. truncatula* inhabited Mt. Elgon [20] and the Kitulo Plateau in southern Tanzania [22]. Both studies, however, used short fragments of the highly conservative nuclear ribosomal 18S gene. Whereas, this genetic marker is sufficient to delimit *Galba* spp. from *Radix natalensis*, it is not suitable for intra-generic studies. Given this situation, it remains currently unclear whether the high-altitude African populations of *Galba* spp. indeed represent *Galba truncatula*. Moreover, it is unknown how these populations are related to populations in Europe, Asia and the Americas. Due to the complete absence of molecular assessments (but see [22]) it is, to date, impossible to reconstruct the centre of origin of *Galba* spp. and to understand when and by what means *Galba* spp. may have colonized Africa and to what extent humans might have been involved in that process.

To shed new light on the phylogeography of *Galba* spp. populations, and its impact on snail-borne

diseases, we examine several African populations using combination of mitochondrial and nuclear DNA markers to determine the identity and biogeographical affinities, reconstruct the colonization history including the likely mode of dispersal, and test for the presence of cryptic *Galba* species in Africa.

Methods

Sampling

The snail specimens studied were collected in Africa between 2010 and 2018. Field trips were conducted in the Atlas Mountains in Morocco, the highlands of Ethiopia, the Eastern Arc Mountains of Tanzania, Mt. Elgon in Uganda and the highlands of Lesotho in southern Africa (Table 1). In addition, material from outside Africa available in the collection of University of Giessen Systematics and Biodiversity (UGSB) was also used. This included material from the type-locality of *G. truncatula* in Thuringia, Germany. Snails were manually collected using a scoop net in stable pools, ponds, marshes, swamps and slow-running waters. Specimens were fixed in 80% ethanol prior to DNA extraction.

DNA extraction, amplification and sequencing

In most cases, DNA was extracted from two *Galba* specimens per locality. DNA extraction from ethanol-preserved snails was performed following the CTAB protocol of [28]. The primers used to amplify a fragment of the *cox1* gene with a target length of 658 bp were LCO1490 and HCO2198 [29]. Amplification of the *LSU* rRNA fragment (*16S*) with a target length of 500 bp was performed with primers 16Sar and 16Sbr [30]. For the nuclear internal transcribed spacer ITS2, primers LT1 and ITS2-RIXO were used [9, 31].

PCR conditions were as described in [32]. Bidirectional sequencing was performed on an ABI 3730 XL sequencer at LGC Genomics, Berlin, Germany. *Galba* spp. samples successfully sequenced comprised two specimens from Germany, three specimens from Greece, two specimens from Slovenia, five specimens from Russia, six specimens from Nepal, one specimen from Ethiopia, five specimens from Lesotho, nine specimens from Morocco, four specimens from Tanzania, and six specimens from Uganda (Table 1).

Phylogenetic analyses

DNA sequences were edited using MEGA v.7.0 [33]. The resulting dataset was complemented with other *Galba* spp. and *Lymnaea* spp. sequences available on GenBank (Table 1). The final dataset comprised a total of 19 specimens. The *16S* partition was aligned using the online program MAFFT [34], whereas Prankster [35] was used to align the ITS2 partition. The final concatenated

alignment was 1494 bp long (*16S*: 434 bp; *cox1*: 655 bp; ITS2: 405 bp). Two outgroups were used for rooting the tree, *Radix natalensis* and *Pseudosuccinea columella* (Table 1).

We used jModelTest v.2.1.4 [36] to identify the best-fit substitution model for running phylogenetic analyses based on Bayesian inference (BI) as implemented in MrBayes v.3.2.6 [37]. Based on the corrected Akaike's information criterion (AICc), the best-fit models were: GTR+ Γ for *16S*, GTR+I+ Γ for *cox1*, and GTR+ Γ for ITS2. We ran two independent Markov Chain Monte Carlo (MCMC) searches (each with four chains) for 1 million generations and sampled every 50th tree and applied a 'burn-in' of 50%. Convergence of the two independent runs was examined *a posteriori* in Tracer 1.5 [38]. Effective sample size (ESS) values of >200 indicated adequate sampling of posteriors distributions. In addition, a maximum likelihood (ML) analysis was conducted using RAxML-HPC2 8.2.10 [39] on the CIPRES Science Gateway [40] by applying the GTR+ Γ model to all partitions and using a stop rule for the bootstrap analysis as recommended.

Estimation of divergence times

Because of the scanty fossil record of *Galba* spp. and lymnaeids in general [4] and given the absence of a specific substitution rate for Lymnaeidae or freshwater pulmonate gastropods in general, we adopted a very conservative approach of dating the molecular phylogeny. We used two substitution rates for *cox1*, i.e. 1%/myr and 2%/myr and estimated divergence times using BEAST v.1.8.4 [41]. Analyses were run for 20 million generations, sampling every 1000th tree. Convergence of runs was analyzed using Tracer v.1.5. Because convergence was not reached and ESS values were < 200, we applied the less complex HKY substitution model to the different partitions (i.e. *16S*: HKY+ Γ ; *cox1*: HKY+I+ Γ ; and ITS: HKY+ Γ). The maximum clade credibility (MCC) tree was identified using TreeAnnotator v.1.8.4 (BEAST package) by applying a 'burn-in' of 50%.

Phylogeographical analyses

Phylogeographical analyses were carried out for the subset of samples from sub-Saharan Africa. The datasets consisted of 11 sequences for *cox1*, 11 sequences for *16S*, and 16 sequences for ITS2 and were individually analyzed. Relationships between haplotypes were calculated using a statistical parsimony network analysis performed using the software tool TCS v.1.21 [42] with a connection limit of 95%. Uncorrected genetic p-distances were calculated in MEGA v.7.0 [33] for within and among major *cox1* clades inferred from the phylogenetic analyses.




Fig. 1 Distribution map of *Galba* in Africa including the sampling for the present study (see Table 1 for details). Four sub-ranges hitherto known of *Galba truncatula* are indicated (adopted from [21] and modified from [24, 26]. Note that occurrences on the Arab Peninsula are not shown here. Black dots denote isolated occurrences; white dots represent subfossil records. Localities of the newly obtained material are shown as coloured stars

Results

Phylogenetic analyses and divergence time estimation

The phylogenetic analyses conducted resulted in a generally highly supported phylogeny (Fig. 2) including a highly supported clade (ML bootstrap values, $bs = 96$; MrBayes posterior probability; $pp = 1.00$, BEAST posterior probability; $bpp = 1.00$) represented by *G. truncatula* comprising samples from Europe (including the type-locality in Thuringia, Germany), Asia, and a single specimen from Morocco. The remaining African samples formed a highly supported monophyletic clade ($bs = 98$; $pp = 1.00$; $bpp = 1.00$) that is referred to as *G. mweruensis* hereafter, which is possibly sister to *G. truncatula* ($bs = 77$, $pp = 0.81$, $bpp = 1.00$). *Galba mweruensis* (Connolly, 1929) is an available name for that clade ([43]; see Discussion). The distinction of *G. mweruensis* from *G. truncatula* is further corroborated by a more comprehensive *cox1*-based phylogeny (Additional file 1: Figure S1) and genetic distances (Table 2). However, both phylogenetic approaches (MrBayes and BEAST) revealed slightly different topologies. According to the MrBayes analysis, a clade of *Lymnaea humilis* and *L. cousini* was sister to the two *Galba* species. They together formed the

sister-group to the remaining South American species (*L. cubensis*, *Lymnaea* sp., and *L. viatrix*). The cryptic species *G. schirazensis* from Iran and *L. diaphana* are more distantly related. In contrast, the BEAST analysis suggests a closer relationship of *G. schirazensis* (Iran) and *Lymnaea* sp. (Colombia) to *L. truncatula* and *L. mweruensis* and also found differences in the more basal phylogenetic relationships.

The split between *G. truncatula* and *G. mweruensis* was estimated to have occurred between $c.3.9$ (95% highest posterior density, 95% HPD: 5.6–10.2) and $c.7.8$ (95% HPD: 2.8–5.1) million years ago (Ma) depending on whether a clock rate of 2%/myr or 1%/myr was used (Additional file 2: Figure S2 and Additional file 3: Figure S3). The diversification of *G. mweruensis* started between $c.1.7$ (95% HPD: 1.1–2.3) and $c.3.4$ (95% HPD: 2.3–4.6) Ma.

Phylogeographical analysis

The *cox1* haplotype network consisted of six haplotypes, two of which belonged to populations from Tanzania and Lesotho each, whereas the single specimens from Ethiopia and Uganda represented unique haplotypes. These geographical haplotypes were all connected except for the populations from Mt. Elgon (Uganda) that were separated by at least 22 mutational steps from the remaining haplotypes and thus represented a distinct haplotype network based on the 95% connection limit (Fig. 3). Similar patterns were also revealed by the 16S and ITS2 datasets. Populations from Tanzania and Ethiopia seem to be more closely related in the two mitochondrial networks, whereas the ITS2 dataset suggested a closer relationship between populations from Ethiopia, Lesotho and Tanzania. The individuals from Mt. Elgon were also not connected with the remaining populations in the 16S network (separated by at least 14 mutational steps) and were separated by 8 mutational steps from the other haplotypes in the ITS2 network based on the 95% connection limit.

The genetic distance within *G. truncatula* was higher (4.4%) than within *G. mweruensis* (1.9%). The uncorrected genetic p-distance between both groups was considerably high (9.0%).

Discussion

Identity of *Galba* in Africa and phylogenetic affinities

This study found two geographically separated species of *Galba* in Africa. *Galba truncatula* is restricted based on the available evidence to the Palaearctic zone of the continent, namely Morocco. All sub-Saharan populations proved to be a distinct species according to the phylogenetic analyses and genetic distance to the sister species

Table 1 Locality, voucher (UGSB no.), and GenBank accession information for the species studied. UGSB is the acronym of the University of Giessen Systematics and Biodiversity collection

| Species | Locality | Latitude | Longitude | Altitude (masl) | Code | UGSB no. | GenBank ID | | |
|---------------------------------|---------------------------------------------------|------------|------------|-----------------|----------|----------|------------|----------|----------|
| | | | | | | | cox1 | 16S | ITS2 |
| <i>Galba mweruensis</i> | Lesotho, Mantsonyane | 29.51682°S | 28.29032°E | 2212 | Gmw15772 | 23470 | MN601402 | MN602685 | MN602657 |
| | | | | | Gmw15773 | 23471 | MN601403 | MN602686 | MN602658 |
| | | | | | Gmw15775 | 23473 | MN601405 | MN602688 | MN602660 |
| | | | | | Gmw15776 | 23474 | MN601406 | MN602689 | MN602661 |
| | Tanzania, Lushoto | 04.44859°S | 38.17837°E | 1639 | Gmw25316 | 20983 | MN601423 | MN602698 | MN602674 |
| | | | | | Gmw25317 | 20984 | MN601424 | MN602699 | MN602675 |
| | | | | | Gmw25318 | 20985 | MN601425 | MN602700 | MN602676 |
| | | | | | Gmw25319 | 20986 | MN601426 | MN602701 | |
| | Ethiopia, Adi Aba Musa, Lake Ashenge | 12.58650°N | 39.52100°E | 2409 | Gmw22773 | 17407 | MN601410 | MN602707 | MN602665 |
| | Uganda, Budadiri, Mt. Elgon, Jackson's Pool | 01.14951°N | 34.51054°E | 3939 | Gmw19054 | 12151 | MN601409 | MN602706 | MN602664 |
| | Uganda, Mt. Elgon | 01.14954°N | 34.54736°E | 3792 | Gmw26767 | 22833 | | | MN602677 |
| | | | | | Gmw26769 | 22835 | | | MN602678 |
| | | | | | Gmw26770 | 22836 | | | MN602679 |
| | | | | | Gmw26771 | 22837 | | | MN602680 |
| | | | | | Gmw26772 | 22838 | | | MN602681 |
| <i>Galba truncatula</i> | Morocco, Marrakech-Safi | 31.15573°N | 07.86678°W | 2100 | Gtr25298 | 18267 | MN601412 | MN602690 | MN602666 |
| | Morocco, Tindighas | 32.68417°N | 05.33972°W | 1982 | Gtr25297 | 18265 | MN601411 | | |
| | Morocco, Marlay youssef Dam | 31.39272°N | 07.15383°W | 167 | Gtr25304 | 20971 | MN601415 | | |
| | | | | | Gtr25305 | 20972 | MN601416 | | |
| | Germany, Thuringia, Ilm River | 50.89112°N | 11.24089°E | 289 | Gtr15785 | 23475 | MN601407 | MN602704 | MN602662 |
| | | | | | Gtr15786 | 23476 | MN601408 | MN602705 | MN602663 |
| | Greece, Rhodos Island, 7 springs dam lake, on mud | 36.25464°N | 28.11596°E | 232 | Gtr25308 | 20975 | MN601419 | MN602694 | MN602670 |
| | | | | | Gtr25306 | 20973 | MN601417 | | |
| | | | | | Gtr25307 | 20974 | MN601418 | MN602693 | MN602669 |
| | Russia, Ilovlya, river near Ilovlya Town | 49.31367°N | 43.97659°E | 43 | Gtr25312 | 20979 | MN601420 | MN602695 | MN602671 |
| | Russia, Moscow Region, Oka River | na | na | | Gtr25313 | 20980 | MN601421 | MN602696 | MN602672 |
| | | | | | Gtr25314 | 20981 | MN601422 | MN602697 | MN602673 |
| | Slovenia, Vrhnik, creek Obrh | 45.69906°N | 14.51176°E | 376 | Gtr25299 | 18543 | MN601413 | MN602691 | MN602667 |
| | | | | | Gtr25301 | 18860 | MN601414 | MN602692 | MN602668 |
| | Nepal, Karnali | 29.26667°N | 82.15933°E | 2300 | Gtr11234 | 23477 | MN601399 | MN602702 | MN602654 |
| | | | | | Gtr12653 | 23478 | | MN602703 | MN602656 |
| <i>Lymnaea schirazensis</i> | Nepal, Bagmati | 29.30000°N | 82.36667°E | 2700 | Gtr11235 | 23479 | MN601400 | MN602684 | MN602655 |
| | Nepal, Bheri | 29.10717°N | 82.58867°E | 2625 | Gtr11237 | 23481 | MN601401 | | |
| | France, Limoges | | | | GB2 | | | HQ283236 | HQ283262 |
| | Iran, Gilan Province, Taleb-Abad River | | | | GB1 | | JF272607 | JF272605 | |
| | | | | | | | | | |
| | USA, New York | | | | GB3 | | | FN182195 | FN182191 |
| | Venezuela, Mucubají | | | | GB4 | | | HQ283237 | HQ283266 |
| | USA, South Carolina | | | | GB5 | | | FN182204 | |
| | Argentina, Lago Escondido | | | | GB6 | | | HQ283241 | HQ283260 |
| | Colombia, Antioquia | | | | GB7 | | | HQ283235 | HQ283263 |
| <i>Lymnaea</i> sp. | Argentina, Rio Negro | | | | GB8 | | | HQ283239 | HQ283265 |
| | | | | | | | | | |
| | | | | | | | | | |
| <i>Radix natalensis</i> | Kenya, Kisumu, Lake Victoria | 00.12739°S | 34.74232°E | 1140 | Rna15771 | 23483 | MN601427 | MN602708 | MN602708 |
| <i>Pseudosuccinea columella</i> | South Africa, Mpumalanga | 24.84539°S | 30.83879°E | 1374 | Pco15787 | 23484 | MN601428 | MN602709 | MN602683 |

Abbreviations: na, not available; masl, meters above sea level

G. truncatula from Europe and Asia. Interestingly, no *G. schirazensis* was found at the examined localities, which further supports the hypothesis that mountain ranges of tropical Africa are inhabited by a species different from *G. truncatula* and its cryptic counterpart *G. schirazensis* has not had opportunity to disperse into these areas or is unable to do so. We therefore propose to use the existing name *G. mweruensis* (Connolly, 1929) for this species that was described based on shell features and size measures (for a comparison of the original type-material and our new populations see Additional file 4: Figure S4; Additional file 5: Table S1). Moreover, it is morphologically indistinguishable from the other two species hitherto known to occur in Africa, i.e. *G. truncatula* and *G. schirazensis* (Additional file 6: Figure S5). *Galba mweruensis* is not the oldest available name for African *Galba* species for which even the section name *Afrogalba* had been introduced by Kruglov & Starobogatov [44]. Another taxon described earlier is *Galba umlaasianus* (Küster, 1862) from the Umlaas River, South Africa. Recent repeated attempts to obtain material from *terra typica* in the Kwa Zulu Natal Province of South Africa unfortunately failed. However, *G. umlaasiana* originally has been referred to as a lowland species of the temperate zones along the coastal regions of South Africa, whereas *G. mweruensis* has been described from mountainous terrain from Mweru town (type-locality) at the foothills of Mt. Kenya, which is somewhat in the core range of the species we found to occur widely in tropical Africa. Attempts to locate a population in the Mweru region in central Kenya in 2010 unfortunately failed. Moreover, Vinarski [45] compared both *G. mweruensis* and *G. umlaasiana* with the newly described *G. robusta* from Yemen and found the former two species to be morphologically different. We therefore propose to use the name *G. mweruensis* for mountainous *Galba* populations until it can be compared with topotypic material of *G. umlaasianus*. The latter taxon might even represent another distinct species given its different altitudinal range and may potentially co-occur with *R. natalensis* in the lower altitudes. Such a co-occurrence has not been observed for *G. mweruensis* in the studies that were conducted in the highlands of Lesotho (as *G. truncatula* in [24]), the Kitulo Plateau in Tanzania [22], and Mt. Elgon in Uganda [20]. In South Africa, however, either *G. truncatula* (*G. umlaasianus*), *L. natalensis* or the invasive *P. columella* have been reported to occur sympatrically [24].

Among the newly genotyped specimens of this species, the population from Mt. Elgon in Uganda is of particular interest. Mandahl-Barth [46] identified a small form of *Galba* at Mt. Elgon at 2770 m and attributed it to *G. mweruensis*. According to the present analyses, this population turned out to be sister to the remaining

populations from Ethiopia, Lesotho and Tanzania, and the Mt. Elgon population was very distantly related to the remaining groups in the phylogeographical analyses. A more detailed analysis that investigates morphological and anatomical characters is needed in order to establish the status of the Mt. Elgon populations compared to their sub-Saharan counterparts. Hubendick [26] had material from the Kenyan slopes of Mt. Elgon and found similarities to *G. truncatula* but treated it as *G. mweruensis*. Isolated records of *Galba* spp. from the eastern part of the DR Congo west of Lake Albert and at Lake Kivu from considerably lower altitudes have not been confirmed during the last decades [21, 47].

The genetic diversity within *G. mweruensis* is comparable to that of other distinct *Galba* species such as *G. schirazensis* [26]. Given the continuous and by far greater distributional range of *G. truncatula*, the higher degree of genetic differentiation in *G. truncatula* compared to *G. mweruensis* is not surprising. Nevertheless, the comparatively high genetic diversity within *G. mweruensis* raises the question as to how this diversity in isolated patches scattered over Africa has evolved and how these areas have been colonized. Further study in detail of several life-history traits for survival in cooler zones could be illuminating.

Colonization history

Our study indicates that subtropical Africa has been colonized only once in either the Pliocene or even Miocene if one considers the age of the most recent common ancestor of *G. truncatula* and *G. mweruensis* as indicative of colonization time. Diversification within the African species *G. mweruensis* is dated to the Plio-Pleistocene and thus human-mediated dispersal can be ruled out for the initial colonization of the mountain ranges. We here applied commonly used substitution rates for mitochondrial markers in invertebrates, i.e. 1%/myr and 2%/myr (i.e. divergence rates of 2%/myr and 4%/myr). Assuming that *Galba* may have evolved with an extremely fast substitution rate of 4%/myr, the split would, of course, become younger (*c.* 2 Ma). However, this would not change our conclusions that the hypothesis of human-mediated dispersal can be rejected. However, the data do not currently allow drawing a final conclusion as to whether Africa has been colonized from Europe, the Near East or South America. The tree topology may favour a colonization scenario out of Europe; however, Asian and especially Near East samples of *G. truncatula* are scarce and *G. robusta* (Yemen) could not be included. Subfossil records in Africa are also not very helpful as they originate from less mountainous regions and are not very informative given the small morphospace occupied by all *Galba* species. However, recent and subfossil





Table 2 Genetic distances of *Galba mweruensis* and *Galba truncatula* based on the *cox1* dataset

| | Uncorrected p-distance (%) | | | K2P model | | |
|---------|----------------------------|----------------------|----------------------------------------------|----------------------|----------------------|----------------------------------------------|
| | <i>G. mweruensis</i> | <i>G. truncatula</i> | <i>G. mweruensis</i> vs <i>G. truncatula</i> | <i>G. mweruensis</i> | <i>G. truncatula</i> | <i>G. mweruensis</i> vs <i>G. truncatula</i> |
| Minimum | 0.2 | 0.0 | 7.1 | — ^a | — ^a | — ^a |
| Maximum | 4.2 | 7.8 | 9.7 | — ^a | — ^a | — ^a |
| Mean | 2.3 | 3.2 | 8.4 | 2.4 | 3.4 | 9.0 |

Note: Uncorrected genetic p-distances and genetic distances based on the K2P model were calculated in MEGA v.7.0 [33]

^a Not calculated



Saharan records [18, 21] may indicate a stepping-stone dispersal for the northern Africa *G. truncatula* populations. The generally much higher lymnaeid diversity in the northern hemisphere makes an ‘out of Africa’ alternative for the *Galba* less likely. However, given the existence of the cryptic *G. schirazensis* in Egypt [27], no conclusion can be drawn here. On the intra-continental scale, a closer relationship between the Northeast and East African populations in comparison to the populations of the highlands of Lesotho would be expected. However, according to our analyses, specimens from Mt. Elgon are genetically more distinct compared to the remaining sub-Saharan haplotypes.

Dispersal by water birds, also at high altitudes, has been commonly shown to be a major factor in range evolution for freshwater molluscs in general [48] and pulmonate snails in particular [49]. To which extent water birds might have been involved in the colonization of these isolated mountain ranges can only be speculated. If such dispersal is as frequent as demonstrated in other regions

[50, 51], *G. mweruensis* should be more widespread across different mountain ranges in sub-Saharan Africa.

Africa has experienced severe climatic fluctuations since the late Miocene and especially in the Plio-Pleistocene [52]. The patchy distribution pattern observed may thus reflect the emergence of climatic refugia in these mountain ranges that acted as islands in the sky [53]. Such relictary species distributions in African mountain ranges have been documented for diverse taxa such as birds [54], flightless insects [55] and frogs [56]. Although the status of *G. umlaasiana* has not been assessed yet, a correlation of cooler climates and the occurrence of *G. mweruensis* is apparent. Alternatively, the presence of the omnipresent and thus potentially competitive *R. natalensis* may considerably restrict the distribution of *G. mweruensis* to more temperate areas. Although mountain ranges are sometimes acting as refugia, they are also sensitive to climate changes [57]. Small and isolated populations might thus go through repeated bottlenecks and might experience local disappearance as found for the *Galba* population on Kitulo Plateau, Tanzania. A recent field survey (FC in October 2018) showed that the swampy habitats where the species earlier occurred [22] had completely dried out. A high estivating potential for *Galba* is, however, reported from highlands of Ethiopia [58].

Parasitological implications of cryptic *Galba* species in Africa

Despite its patchy continental distribution, *G. mweruensis* is well established, especially in the extensive sub-ranges (Fig. 1). We here confirmed its presence in regions where it has not been observed for decades such as the Usambara Mountains (Tanzania) or Mt. Elgon in Uganda. It is also the predominant snail species in the highlands of Ethiopia and Lesotho and thus should be the intermediate host for livestock fascioliasis and potentially other trematode infections in that region [19, 59]. Dinnik & Dinnik [60] already pointed out that *G. mweruensis* is the intermediate host of both liver flukes, *F. hepatica* and *F. gigantica*, and thus not only represent major threats for livestock. For livestock, considerable economic losses are known from several African countries [61]. We suggest that there is a need to now ascertain the level of snail-parasite compatibility of *G. mweruensis* with several isolates of *F. hepatica* and *F. gigantica*, especially where these snails are found in cattle farmed areas.

Although estimating the prevalence of human fascioliasis is challenging [62], infection risks should be considered high wherever the intermediate host occurs [22]. Outbreaks can happen quickly, and the extent is often

underestimated as recently outlined for the mountains in northern Tanzania [63]. Unlike with other human snail-borne diseases such as schistosomiasis, there is a high prevalence in high altitude regions. A prime example is the endemic in the Andean Altiplano [14, 64]. Although high mountainous regions are still considerably remote and less densely populated in Africa, there is a growing demand for land and thus humans increasingly occupying high elevations [65]. Even touristic activities such as trekking and mountain climbing are on the rise in basically all the mountain ranges where *G. mweruensis* occurs so further surveillance is warranted. Therefore, more dedicated surveys on infection and prevalence rates and the study of parasites actually hosted by *G. mweruensis* are necessary in all the areas where this species is established [20]. Whereas *G. schirazensis* is not particularly involved in transmission of *F. hepatica* [27], high rates of infection have been reported for *G. mweruensis* (originally *G. truncatula*) populations from Lesotho and Ethiopia [58, 66].

Conclusions

This study has identified a hitherto neglected distinct species, *G. mweruensis*, as a host of *F. hepatica* throughout sub-Saharan Africa. It had previously been considered to be conspecific with Eurasian *G. truncatula*, a well-known and globally intermediate host species for several trematode parasites. Following our findings, a closer examination of the parasite communities hosted by *G. mweruensis* is needed in order to understand transmission patterns in highlands throughout eastern and southern Africa. Other high altitudes areas in Africa are to be surveyed for this species and veterinary and human health concerns have to be evaluated under the new precondition. It would be also interesting to study host specificity and potential climatic adaptations of both the host and the preferred temperature range of *F. hepatica* in Africa. The nature of striking non-overlap in occurrences between the omnipresent *R. natalensis* and *G. mweruensis* deserves more scientific attention because of its evolutionary implications and possible epidemiological cross-over as implicated host of *F. gigantica* and *F. hepatica*.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13071-019-3825-9>.

Additional file 1: Figure S1. Bayesian inference phylogram based on *cox1*. The two outgroups have been removed *a posteriori*. Bayesian posterior probabilities are provided next to each node (top: MrBayes, bottom: RAxML). Sequences obtained from GenBank are labelled plain whereas new sequences from this study are bold. Nodes 1 and 2 indicate the nodes for which divergence time estimates are discussed.

Additional file 2: Figure S2. BEAST molecular clock tree based on an HKY model and a substitution rate of 1%.

Additional file 3: Figure S3. BEAST molecular clock tree based on an HKY model and a substitution rate of 2%.

Additional file 4: Figure S4. Shell measurements of *Galba mweruensis* populations in comparison to the type specimen as described in Connolly, 1929 (p. 175).

Additional file 5: Table S1. Shell measurements of *Galba mweruensis* in the highlands of Lesotho, Tanzania and Mt. Elgon in Uganda.

Additional file 6: Figure S5. Shell, soft body anatomy and reproductive organs of *Galba mweruensis* from Lesotho (Mantsonyane). **Abbreviations:** BC, bursa copulatrix; PHT, phallotheca; PRP, praeputium; VD, vas deferens.

Abbreviations

asl: above sea level; ESS: effective sample size; Gtr: *Galba truncatula*; Gmw: *Galba mweruensis*; Pco: *Pseudosuccinea columella*; PCR: polymerase chain reaction; Rna: *Radix natalensis*; UGSB: University of Giessen Systematics and Biodiversity collection.

Acknowledgements

We thank D. Engelhard (Giessen) and T. Geertz (Radolfzell) for collecting on the top of Mt. Elgon, Uganda. We further gratefully acknowledge material provided by K. Boulafasser (Casablanca), U. Bößneck (Erfurt), D. Delicado (Giessen), T. Hauße (Giessen), A. Kossler (Berlin), and M. Vinarski (St. Petersburg). We thank P. Glöer (Hettingen) for his help with the dissections. The comments of two anonymous reviewers helped to improve a previous version of the manuscript and are gratefully acknowledged.

Authors' contributions

AM and CA conceived the study. AM produced the sequences and performed data analyses, with the help of CC and BS. CA, CC and FC collected part of the material, and all authors were involved in data interpretation. AM produced the figures. All authors critically reviewed the manuscript. All authors read and approved the final manuscript.

Funding

AM received a PhD scholarship from the German Academic Exchange Service (DAAD), BS was supported by a Deutsche Forschungsgemeinschaft grant (DFG STE 2460/2-1) and FC was supported by a postdoctoral fellowship of the Humboldt-Foundation. CA received support from the Deutsche Forschungsgemeinschaft (DFG AL 1076/5-2, AL 1076/6-2).

Availability of data and materials

All data generated or analysed during this study are included in the article and its additional files. The newly generated sequences were submitted to the NCBI GenBank database under the accession numbers MN601399–MN601428 for *cox1*, MN602684–MN602709 for *16S*, and MN602654–MN602683 for ITS2.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ Department of Animal Ecology and Systematics, Justus Liebig University Giessen, Giessen, Germany. ² Department of Environmental Sciences, Zoological Institute, University of Basel, Basel, Switzerland. ³ Department of Zoology, University of Dar es Salaam, Dar es Salaam, Tanzania. ⁴ Department of Tropical Disease Biology, Liverpool School of Tropical Medicine, Liverpool, UK. ⁵ Department of Biology, Mbarara University of Science and Technology, Mbarara, Uganda.

Received: 30 April 2019 Accepted: 23 November 2019
Published online: 04 December 2019

References

- Mas-Coma S, Funatsu IR, Bargues MD. *Fasciola hepatica* and lymnaeid snails occurring at very high altitude in South America. *Parasitology*. 2001;123:115–27.
- Lu XT, Gu QY, Limpanont Y, Song LG, Wu ZD, Okanurak K, Lv ZY. Snail-borne parasitic diseases: an update on global epidemiological distribution, transmission interruption and control methods. *Infect Dis Poverty*. 2018;9(7):28.
- Correa AC, Escobar JS, Durand P, Renaud F, David P, Jarne P, et al. Bridging gaps in the molecular phylogeny of the Lymnaeidae (Gastropoda: Pulmonata), vectors of fascioliasis. *BMC Evol Biol*. 2010;10:381.
- Vinarski MV, Clewing C, Albrecht C. Lymnaeidae Rafinesque, 1815. In: Lydeard C, Cummings KS, editors. *Freshwater mollusks of the world: a distribution atlas*. Baltimore: JHU Press; 2019. p. 158–62.
- Dillon RT. *The ecology of freshwater molluscs*. Cambridge: Cambridge University Press; 2000.
- Mas-Coma S, Valero MA, Bargues MD. Fascioliasis. Digenetic trematodes. New York: Springer; 2014. p. 77–114.
- Bölsneck U. Leben am Limit: Besiedlung von Süßwasser-Habitaten extremer Hochlagen Asiens, Amerikas und Afrikas durch Mollusken (Mollusca: Bivalvia & Gastropoda). In: Hartmann M, Weipert J, editors. *Biodiversität und Naturlandschaft im Himalaya IV*. Erfurt: Selfpublished; 2012. p. 103–6.
- Lydeard C, Cummings KS. *Freshwater mollusks of the world: a distribution atlas*. Baltimore: JHU Press; 2019.
- Bargues MD, Vigo M, Horak P, Dvorak J, Patzner RA, Pointier JP, et al. European Lymnaeidae (Mollusca: Gastropoda), intermediate hosts of trematodiasis, based on nuclear ribosomal DNA ITS-2 sequences. *Infect Genet Evol*. 2001;1:85–107.
- Remigio EA, Blair D. Molecular systematics of the freshwater snail family Lymnaeidae (Pulmonata: Basommatophora) utilising mitochondrial ribosomal DNA sequences. *J Molluscan Stud*. 1997;63:173–85.
- Puslednik L, Ponder WF, Dowton M, Davis AR. Examining the phylogeny of the Australasian Lymnaeidae (Heterobranchia: Pulmonata: Gastropoda) using mitochondrial, nuclear and morphological markers. *Mol Phylog Evol*. 2009;52:643–59.
- Glöer P. Die Süßwassergastropoden Nord-und Mitteleuropas. Die Tierwelt Deutschlands 73. Hackenheim: ConchBooks; 2002.
- Seddon MB, Kebapçı U, Van Damme D. *Galba truncatula*. The IUCN Red List of Threatened Species 2015. <http://www.iucnredlist.org/details/10.2305>. Accessed 19 Apr 2019.
- Bargues MD, Artigas P, Khoubbane M, Ortiz P, Naquira C, Mas-Coma S. Molecular characterisation of *Galba truncatula*, *Lymnaea neotropica* and *L. schirazensis* from Cajamarca, Peru and their potential role in transmission of human and animal fascioliasis. *Parasit Vectors*. 2012;5:174.
- Alda P, Lounnas M, Vázquez AA, Ayaqui R, Calvopiña M, Celi-Erazo M, et al. A new multiplex PCR assay to distinguish among three cryptic *Galba* species, intermediate hosts of *Fasciola hepatica*. *Vet Parasitol*. 2018;251:101–5.
- Duffy T, Kleiman P, Pietrovsky S, Issia L, Schijman AG, Wisnivesky-Colli C. Real-time PCR strategy for rapid discrimination among main lymnaeid species from Argentina. *Acta Trop*. 2009;109:1–4.
- Trouvé S, Degen L, Meunier C, Tirard C, Hurtrez-Boussès S, Durand P, et al. Microsatellites in the hermaphroditic snail, *Lymnaea truncatula*, intermediate host of the liver fluke, *Fasciola hepatica*. *Mol Ecol*. 2000;9:1662–4.
- Van Damme D. *The freshwater Mollusca of northern Africa: distribution, biogeography and palaeoecology*. The Hague: Dr. W. Junk Publishers; 1984.
- Goll PH, Scott JM. Fascioliasis in the Ethiopian central highlands. Dynamics of intermediate snail host [*Lymnaea truncatula*] populations and their relation to infection in sheep [*Fasciola hepatica*, liver fluke]. *Misc Rep-C Overseas Pest Res UK*. 1979;47:1–12.
- Howell A, Mugisha L, Davies J, LaCourse EJ, Claridge J, Williams DJ, et al. Bovine fasciolosis at increasing altitudes: parasitological and malacological sampling on the slopes of Mount Elgon, Uganda. *Parasit Vectors*. 2012;5:196.
- Brown DS. *Freshwater snails of Africa and their medical importance*. London: CRC Press; 1994.
- Walker SM, Makundi AE, Namuba FV, Kassuku AA, Keyyu J, Hoey EM, et al. The distribution of *Fasciola hepatica* and *Fasciola gigantica* within southern Tanzania—constraints associated with the intermediate host. *Parasitology*. 2008;135:495–503.
- De Kock KN, Wolmarans CT, Bornmas M. Distribution and habitats of the snail *Lymnaea truncatula*, intermediate host of the liver fluke *Fasciola hepatica*, in South Africa. *J S Afr Vet Ass*. 2003;74:117–22.
- De Kock KN, Wolmarans CT. The geographical distribution and habitats of three liver fluke intermediate hosts in South Africa and the health implications involved. *Suid-Afrikaanse Tydskrif vir Natuurwetenskap en Tegnologie*. 2008;27:1–16.
- Mas-Coma S, Valero MA, Bargues MD. *Fasciola*, lymnaeids and human fascioliasis, with a global overview on disease transmission, epidemiology, evolutionary genetics, molecular epidemiology and control. *Adv Parasitol*. 2009;69:41–146.
- Hubendick B. *Recent Lymnaeidae: their variation, morphology, taxonomy, nomenclature, and distribution*. Berlin: Almqvist & Wiksell; 1951.
- Bargues MD, Artigas P, Khoubbane M, Flores R, Glöer P, Rojas-García R, et al. *Lymnaea schirazensis*, an overlooked snail distorting fascioliasis data: genotype, phenotype, ecology, worldwide spread, susceptibility, applicability. *PLoS ONE*. 2011;6:24567.
- Wilke T, Davis GM, Qiu DC, Spear RC. Extreme mitochondrial sequence diversity in the intermediate schistosomiasis host *Oncomelania hupensis robertsoni*: another case of ancestral polymorphism? *Malacologia*. 2006;48:143–57.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol*. 1994;5:294–9.
- Palumbi S. Simple fool's guide to PCR. Honolulu: Department of Zoology and Kewalo Marine Laboratory; 1991.
- Almeyda-Artigas RJ, Bargues MD, Mas-Coma S. ITS-2 rDNA sequencing of *Gnathostoma* species (Nematoda) and elucidation of the species causing human gnathostomiasis in the Americas. *J Parasitol*. 2000;86:537–44.
- Stelbrink B, Shirokaya AA, Föller K, Wilke T, Albrecht C. Origin and diversification of Lake Ohrid's endemic acroloxid limpets: the role of geography and ecology. *BMC Evol Biol*. 2016;16:273.
- Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol Biol Evol*. 2016;33:1870–4.
- Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol*. 2013;30:772–80.
- Löytynoja A, Goldman N. web PRANK: a phylogeny-aware multiple sequence aligner with interactive alignment browser. *BMC Bioinform*. 2010;11:579.
- Posada D, Crandall KA. Modeltest: testing the model of DNA substitution. *Bioinformatics*. 1998;14:817–8.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, et al. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol*. 2012;61:539–42.
- Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol Biol*. 2007. p. 214.
- Stamatakis A. RAXML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. 2014;30:1312–3.
- Miller MA, Pfeiffer W, Schwartz T. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. 2010 Gateway Computing Environments Workshop (GCE); 2010. p. 1–8.
- Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Mol Biol Evol*. 2012;29:1969–73.
- Clement M, Posada D, Crandall KA. TCS: a computer program to estimate gene genealogies. *Mol Ecol*. 2000;9:1657–9.
- Connolly M. Notes on African non-marine Mollusca, with descriptions of many new species. *Ann Mag Nat Hist*. 1929;156:78.
- Kruglov ND, Starobogatov YI. The volume of the subgenus *Galba* and of the other *Lymnaea* (Gastropoda, Pulmonata) subgenera, similar to it. *Zool Zhurnal*. 1985;64:24–35.
- Vinarski MV. *Galba robusta* sp. nov. from Yemen (Gastropoda: Lymnaeidae). *Zoosyst Rossica*. 2018;27:3–10.
- Mandahl-Barth G. *The freshwater mollusks of Uganda and adjacent territories*. Tervuren: Royal Museum of Belgian Congo; 1954.
- Fain A. Ecological and parasitological notes on *L. truncatula* in the Belgian Congo. *Ann Soc Belge Med Trop*. 1951;31:149–52.

48. Clewing C, Bössneck U, von Oheimb PV, Albrecht C. Molecular phylogeny and biogeography of a high mountain bivalve fauna: the Sphaeriidae of the Tibetan Plateau. *Malacologia*. 2013;56:231–53.
49. Von Oheimb PV, Albrecht C, Riedel F, Bössneck U, Zhang H, Wilke T. Testing the role of the Himalaya Mountains as a dispersal barrier in freshwater gastropods (*Gyraulus* spp.). *Biol J Linn Soc*. 2013;109:526–34.
50. Green AJ, Figuerola J. Recent advances in the study of long-distance dispersal of aquatic invertebrates *via* birds. *Diver Distr*. 2005;11:149–56.
51. Kappes H, Haase P. Slow, but steady: dispersal of freshwater molluscs. *Aquat Sci*. 2012;74:1–14.
52. Cox SC, Prys-Jones RP, Habel JC, Amakobe BA, Day JJ. Niche divergence promotes rapid diversification of East African sky island white-eyes (Aves: Zosteropidae). *Mol Ecol*. 2014;23:4103–18.
53. Ring U, Albrecht C, Schrenk F. The East African Rift system: tectonics, climate and biodiversity. In: Hoorn C, Perrigo A, Antonelly A, editors. *Mountains, climate and biodiversity*. Oxford: Wiley Blackwell; 2018. p. 391–411.
54. McCormack JE, Huang H, Knowles LL, Gillespie R, Clague D. Sky islands. *Encycl Isl*. 2009;4:841–3.
55. Brühl C. Flightless insects: a test case for historical relationships of African mountains. *J Biogeogr*. 1997;24:233–50.
56. Lawson LP. The discordance of diversification: evolution in the tropical montane frogs of the Eastern Arc Mountains of Tanzania. *Mol Ecol*. 2010;19:4046–60.
57. Peters MK, Hemp A, Appelhaus T, Becker JN, Behler C, Classen A, et al. Climate-land-use interactions shape tropical mountain biodiversity and ecosystem functions. *Nature*. 2019;568:7750.
58. Goll PH, Scott JM. The interrelationship of *Lymnaea truncatula* and ovine fascioliasis in the Ethiopian central highlands. *Br Vet J*. 1978;134:551–5.
59. Prinsloo JF, Van Eeden JA. The distribution of the freshwater molluscs in Lesotho with particular reference to the intermediate host of *Fasciola hepatica*. *Wet Byd Potchefstroomse Univ, B, Natuurwetenskappe*. 1973;57:1–11.
60. Dinnik JA, Dinnik NN. A mud snail, *Lymnaea mweruensis* Connolly as an intermediate host of both liver flukes *Fasciola hepatica* L. and *Fasciola gigantica* Cobbold. *Rep E Afr Vet Res Org*. 1957;1:50–2.
61. Mehmood K, Zhang H, Sabir AJ, Abbas RZ, Ijaz M, Durrani AZ, et al. A review on epidemiology, global prevalence and economical losses of fasciolosis in ruminants. *Microb Pathogenesis*. 2017;109:253–62.
62. Mas-Coma S, Bargues MD, Valero MA. Human fascioliasis infection sources, their diversity, incidence factors, analytical methods and prevention measures. *Parasitology*. 2018;145:1665–99.
63. Abdul-Hamid L, Mchale DN, Nyindo M. Diagnosis of human fascioliasis in Arusha region, northern Tanzania by microscopy and clinical manifestations. *Tanzania Vet J*. 2017;32:55–61.
64. Standley CJ, Prepelitchi L, Pietrovsky SM, Issia L, Stothard JR, Wisnivesky-Colli C. Molecular characterization of cryptic and sympatric lymnaeid species from the *Galba/Fossaria* group in Mendoza Province, Northern Patagonia, Argentina. *Parasit Vectors*. 2013;6:304.
65. Kilungu H, Leemans R, Munishi PK, Nicholls S, Amelung B. Forty years of climate and land-cover change and its effects on tourism resources in Kilimanjaro National Park. *Tour Plan Development*. 2019;16:235–53.
66. Prinsloo JF, Van Eeden JA. Population dynamics of freshwater snails in Lesotho with particular reference to *Lymnaea truncatula*: the intermediate host of *Fasciola hepatica*. *Wet Byd Potchefstroomse Univ, B, Natuurwetenskappe*. 1976;72:1–60.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

